



RayBiotech, Inc.

3607 Parkway Lane suite 200
Norcross, GA 30092
Tel: 770-729-2992, 1-888-494-8555
Fax: 770-206-2393
Website: www.raybiotech.com
Email: info@raybiotech.com

Certificate of Analysis and Data Sheet

Human Factor IX

Catalog No.

228-10434

Source:

Human Plasma.

Synonyms:

Coagulation factor IX, EC 3.4.21.22, Christmas factor, Plasma thromboplastin component, PTC, F9, FIX, HEMB, MGC129641, MGC129642, GLA domain, Factor IX.

Introduction:

Human Factor IX also called Christmas-Factor is a glycoprotein, which is synthesized in the liver and belongs to the serine proteases system and is part of the S1 peptidase family.

Lack of Factor-IX causes Hemophilia-B meaning Christmas Disease. Factor-IX has a N-terminus region which contains 12xGla residues which assist the calcium dependant binding of Factor-IX to the phospholipid surface. Factor-IX is activated by either factor XIa or the factor VIIa/tissue factor/phospholipid complex. Cleavage yields the intermediate IXa, which is subsequently converted to the fully active form IXab.

Factor-IX binds initially to exosites on the factor XIa heavy chain, followed by interaction at the active site with subsequent bond cleavage. Coagulation factor IX is activated by interaction with the erythrocyte membrane, causing intrinsic coagulation. Chaperones & lectins act simultaneously to guarantee the proper folding of Factor-IX and the retention of mutant molecules. Human Factor IX, activated by either the Contact or Tissue Factor Pathway, is responsible for the activation of Factor X to Xa.

Description:

Human Factor-IX produced from fresh frozen human plasma is a glycosylated polypeptide chain having a molecular mass of 56 kDa.

Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.


Formulation

The Factor-IX was lyophilized from a sterile solution containing 20mM Tris-HCl pH-7.4, 0.1M NaCl and 1mM Benzamidine.

Solubility

It is recommended to reconstitute the lyophilized 100U Factor-IX in sterile 100μl of 18MΩ-cm H₂O, which can then be further diluted to other aqueous solutions.

**The products are furnished for LABORATORY RESEARCH USE ONLY.
Not for diagnostic or therapeutic use.**



RayBiotech, Inc.

3607 Parkway Lane suite 200
Norcross, GA 30092
Tel: 770-729-2992, 1-888-494-8555
Fax: 770-206-2393
Website: www.raybiotech.com
Email: info@raybiotech.com

Purity:

Greater than 95.0% as determined by SDS-PAGE.

Stability:

Lyophilized Factor-IX although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Factor-IX should be stored at 4°C between 2-7 days and for future use below -18°C.

Please prevent freeze-thaw cycles.

Human Virus Test

Human plasma was tested and found negative for HIV-1, HIV-2, Hepatitis B Surface antigen and HCV. Donors are screened for CJD (Creutzfeldt-Jakob Disease).

Biological Activity

The activity per mg was tested and found to be 195 Units per mg.

**The products are furnished for LABORATORY RESEARCH USE ONLY.
Not for diagnostic or therapeutic use.**