

# Fau I

CCCGCNNNN<sup>^</sup>

## AccuCut Restriction Endonuclease

- Cat. No. E-1661 50 Units  
E-1662 250 Units

- Lot No. : 02C151491H8A3

- Supplied with Enzyme

10X AccuCut™*greeN* Buffer : 1 mL  
100 mM pH 7.6 Tris-HCl  
100 mM MgCl<sub>2</sub>  
10 mM DTT

- Store at -20 °C.

- Source : *Flavobacterium aquatili*.

- Concentration : 1 Units/μL

- Reaction Condition

- 10X AccuCut™ *greeN* Buffer  
- Incubate at 55 °C.

- Storage Buffer

20 mM pH 7.5, Tris-HCl  
50 mM KCl  
1 mM EDTA  
10 mM 2-mercaptoethanol  
50% Glycerol

- Heat inactivation : 65 °C for 20 minutes.

• **Unit definition** : One unit of restriction endonuclease activity is defined as the amount of enzyme required to completely digest 1 μg of substrate DNA in a total reaction volume of 50 μL in one hour using the AccuCut™ buffer provided. Incubations are performed in 1.5 mL tubes at the appropriate incubation temperature as indicated in the Product Profile.

- **Isoschizomer** : None.

- **Neoschizomer** : Unfound

• **Reactivity on methylated substrate DNA**: Unidentified

- **Ref**) 1. *IN<8>Dedkov, V.S., Repin, V.E., Rechkunova, N.I., Degtyarev, S.K., Verhosina, V.A., Vinogradova, T.P., (1990) Izv. Sib. Otd. Akad. Nauk SSSR, vol. 1, pp. 35-37.*  
2. *Degtyarev, S.K., Kolyhalov, A.A., Rechkunova, N.I., Dedkov, V.S., (1989) Bioorg. Khim., vol. 15, pp. 130-132.*  
3. *Degtyarev, S.K., Netesova, N.A., Chizikov, V.E., Abdurashitov, M.A., Unpublished observations.*

## Quality Control

- **Overdigestion Assay** :

No nonspecific activity was detected after incubation of 1 μg of λ DNA with 50 units of *Fau I* for 15 hours.

\* Conditions of low ionic strength, high enzyme concentration, glycerol concentration >5%, or pH >8.0 may result in star activity.

- **Nuclease Contamination Assay** :

No altered pattern was detected after incubation of 1 μg of substrate DNA with *Fau I* in 50 μL reaction volume with the supplied AccuCut™ buffer overnight.

- **Ligation and Recutting Assay** :

This assay is used to test for exonuclease activity that would degrade the termini of restriction fragments, resulting in inhibition of ligation and of subsequent digestion of ligated fragments. After 40-fold overdigestion with *Fau I*, 95% of the DNA fragments can be ligated and recut with *Fau I*.