MONOCLONAL ANTIBODY



## **Anti-Peroxiredoxin III (4G10)**

**Background**: Peroxiredoxin (Prx) is a peroxidase growing family, mammalian members have been known to with cell proliferation, differentiation, and apoptosis.

Many isoforms (about 50 proteins), collected in accordance to the amino acid sequence particularly amino-terminal region containing active site cysteine residue, and the thiol-specific antioxidant activity, distribute throughout all the kingdoms. Among them, mammalian Prx consists of 6 different members grouped into typical 2-Cys, atypical 2-Cys Prx, and 1-Cys Prx. Except Prx VI belonging to 1-Cys Prx subgroup, the other five 2-Cys Prx isotypes have the thioredoxin-dependent peroxidase activity utilizing thioredoxin, thioredoxin reductase, and NADPH as a reducing system. Mammalian Prxs are 20 -30 kilodalton in molecular size and vary in subcellular localization: Prx I, II, and VI in cytosol, Prx III in mitochondria, Prx IV in ER and secretion, Prx V showing complicated distribution including peroxisome, mitochondria and cytosol (1).

**Immunogen :** Recombinant human protein purified from *E.coli* 

**Host:** Mouse

**Size**: 100ul


**Isotype**: IgG1, k Clone number: 4G10


**Composition :** PBS containing 50% glycerol

**Positive control :** HeLa cell lysates

**Storage :** Store for 1 year at -20°C from date

of shipment





## **Applications:**

**ELISA** 

Western Blotting (1:1000)

Immunoprecipitation(1-2ul/400ul cell lysates)

## **Background Reference:**

1) Wood, Z. A. et al. (2003) Trends Biochem Sci. 28(1):32-40.

FOR RESEARCH PURPOSE ONLY NOT FOR DIAGNOSTIC OR THERAPEUTIC USE