www.biossusa.com support@biossusa.com 800.501.7654 [DOMESTIC] +1.781.569.5821 [INTERNATIONAL]

Bioss

bs-6982R

Rabbit Anti-Neutrophil Elastase/ELANE Polyclonal Antibody

Primary Antibodies

Background:

Elastases form a subfamily of serine proteases that hydrolyze many proteins in addition to elastin. Humans have six elastase genes which encode the structurally similar proteins. The product of this gene hydrolyzes proteins within specialized neutrophil lysosomes, called azurophil granules, as well as proteins of the extracellular matrix following the protein's release from activated neutrophils. The enzyme may play a role in degenerative and inflammatory diseases by its proteolysis of collagen-IV and elastin of the extracellular matrix. This protein degrades the outer membrane protein A (OmpA) of E. coli as well as the virulence factors of such bacteria as Shigella, Salmonella and Yersinia. Mutations in this gene are associated with cyclic neutropenia and severe congenital neutropenia (SCN). This gene is clustered with other serine protease gene family members, azurocidin 1 and proteinase 3 genes, at chromosome 19pter. All 3 genes are expressed coordinately and their protein products are packaged together into azurophil granules during neutrophil differentiation. [provided by RefSeq, May 2009].

Source/Purification:

KLH conjugated synthetic peptide derived from human Neutrophil Elastase/ELANE. Was purified by Protein A and peptide affinity chromatography.

 $\textbf{Storage:} \ \ \text{Prepared as lyophilized powder or liquid and shipped on ice. Store at -20°C for one year.}$

Reconstitution:

If the antibody is in liquid form, no reconstitution needed.

Reconstitution is only required for the lyophilized antibody. Please refer to the reconstitution instruction card in the package.

For full size images and description please click HERE.

Size: 100ul or 100ug lyophilized

Concentration: 1ug/uL

Host: Rabbit

Reactivities: Human, Mouse, Rat, Dog,

Application:

WB(1:100-500)ELISA(1:500-1000)IHC-P(1:100-500)IHC-F(1:100-500)

• FACS(1:20-100)

• IF(1:50-200)

 Not yet tested in other applications.
Optimal working dilutions must be determined by the end user.

Antibody Type: Polyclonal

Isotype: IgG

Molecular Weight: 26kDa

Preservatives:

10ug/uL BSA and 0.1% NaN3.

For research use only. CAUTION: Not for human or animal therapeutic or diagnostic use.